

Corning® Extracellular Matrix Proteins

Biological and synthetic surfaces for 2D and 3D cell culture

Providing a wide range of animal, human, and synthetic ECMs for researchers to support improved cell attachment, propagation, differentiation, and migration. Our extensive experience in protein purification along with rigorous quality assurance testing guarantees high-quality consistent products.



Features

- Provides a wide range of animal, human, and synthetic products to meet the needs of many cell culture systems
- Enables 3D cell culture with gels made from reconstituted basement membranes, purified proteins, or synthetic peptide hydrogel
- Supports in vivo studies, such as cell engraftment, with Corning Matrigel® matrix and Corning PuraMatrix™ peptide hydrogel
- Ensures high-quality products with extensive quality control testing
- Saves human embryonic stem cell researchers time when using Corning Matrigel hESC-qualified matrix

Attachment and differentiation of normal and transformed cells

Corning Matrigel matrix, certified LDEV-free and the trusted leading basement membrane, provides a physiologically relevant environment for studies of cell morphology, biochemical function, cell migration or invasion, and gene expression. Corning Matrigel matrix is effective for the attachment and differentiation of both normal and transformed anchorage dependent epitheloid cells as well as other cell types including neurons, Sertoli cells, chick lens, vascular endothelial cells, and hepatocytes.

Variety of purified and synthetic proteins and attachment factors

For more defined culture systems, Corning provides a wide range of purified proteins for cell attachment including collagen types I to VI, laminin, fibronectin, osteopontin, and vitronectin. For studies of human cells that require **xeno-free** culture conditions, Corning offers human collagens, fibronectin, osteopontin, and vitronectin. For researchers who need **animal-free** culture systems, we offer two synthetic products Poly-D-Lysine for 2D cell culture and Corning PuraMatrix peptide hydrogel for 3D cell culture. Poly-D-Lysine promotes cell attachment and/or differentiation for a variety of cell types including transfected cell lines, neuronal cell lines, glial cells, and primary neurons.

Wide range of applications and cited references

Corning ECMs are used for a range applications and cell types including in vitro and in vivo angiogenesis, cell migration and invasion, three-dimensional cell culture, neuronal cell culture, primary hepatocyte culture, culturing human embryonic stem (hES), and induced pluripotent stem (iPS) cells. Be confident in your results with our ECMs which have been cited in more than 5,400 scientific articles (see the following table for examples of these citations).

CORNING

Corning® ECM Product Reference Guide

Product	Chemotaxis/Migration	Invasion	3D Culture	Neuronal Cells	Hepatocytes	Endothelial Cells	Tumor Cells	Muscle Cells	Epithelial Cells	Osteoclasts	hES and iPS cells	Non-specific Cell Attachment
Corning Cell-Tak™ cell and tissue adhesive Supports surface receptor-independent cell attachment. It can simplify the manipulation of weakly adherent cells and tissue sections in immunofluorescence, in situ hybridization, and immunohistochemistry assays.	–	–	25	–	–	50	–	–	–	–	–	80, 81, 82, 83, 84, 85
Corning Matrigel® matrix Growth Factor Reduced is suited for applications where a more highly defined basement membrane preparation is desired. Phenol Red-Free is recommended for assays which require color detection. High Concentration is suited for in vivo applications where a high protein concentration augments growth of tumors. hESC-qualified Matrix supports pluripotent human embryonic stem cell (hESC) growth with mTeSR®1 media from STEMCELL Technologies.	–	11, 14	19	–	40, 42, 43, 44, 46	49, 52, 53, 54, 55, 57	11, 19, 63, 66, 67, 68	–	19, 63, 66, 70	–	73, 74, 75, 76, 77, 78, 79	–
Collagen Has been shown to be involved in cell attachment and differentiation, integrin binding, and chemotaxis.	1, 2, 5	11, 13	16, 17, 18	28, 33	41, 45, 47, 48	51, 56, 58, 60	1, 69	–	–	72	–	17

References

- Hall, CL, et al. *Cancer Res.* 66:8648 (2006).
- Xue, C, et al. *Cancer Res.* 66:192 (2006).
- Azare, J, et al. *Mol. Cell Biol.* 27:4444 (2007).
- Brideau, G, et al. *Cancer Res.* 67:11528 (2007).
- Goldman, J, et al. *FASEB J.* 21:1003 (2007).
- Weaver, AK, et al. *J. Biol. Chem.* 282:31558 (2007).
- Komander, D, et al. *Mol. Biol. Cell.* 19:4837 (2008).
- Tuinstra, RA, et al. *Proc. Natl. Acad. Sci.* 105:5057 (2008).
- Chen, RQ, et al. *J. Biol. Chem.* 284:16752 (2009).
- Zhao, Y, et al. *J. Biol. Chem.* 284:1385 (2009).
- Horak, CE, et al. *Cancer Res.* 67:11751 (2007).
- Biname, F, et al. *Mol. Biol. Cell.* 19:945 (2008).
- Mack, PJ, et al. *J. Biol. Chem.* 284:8412 (2009).
- Yang, J, et al. *Proc. Natl. Acad. Sci.* 106:3913 (2009).
- Semino, CE, et al. *Differentiation.* 71:262 (2003).
- Guo, C & Piacentini, L. *J. Biol. Chem.* 278:46699 (2003).
- Soundararajan, P, et al. *J. Neurosci.* 26:3526 (2006).
- Chen, T, et al. *Stem Cells.* 25:392 (2007).
- Lee, GY, et al. *Nat. Methods.* 4:359 (2007).
- Wang, S, et al. *Tissue Eng.* 14:227 (2008).
- Aguirre, A, et al. *J. Neurosci.* 25:11092 (2005).
- Flanagan, LA, et al. *J. Neurosci. Res.* 83:845 (2006).
- Gelain, F, et al. *PLoS ONE.* 1:e119 (2006).
- Kim, WY, et al. *Neuron.* 52:981 (2006).
- Risner, JR & Holt JR. *J. Neurophysiol.* 96:2364 (2006).
- Biernaskie, J, et al. *J. Neurosci.* 27:9545 (2007).
- Hall, BJ, et al. *J. Neurosci.* 27:13446 (2007).
- Iacovitti, L, et al. *Brain Res.* 1127:19 (2007).
- Liew, CG, et al. *Stem Cells.* 25:1521 (2007).
- Shrestha, B & Diamond, MS. *J. Virol.* 81:11749 (2007).
- Tanaka, S, et al. *J. Biol. Chem.* 282:10506 (2007).
- Bard, L, et al. *J. Neurosci.* 28:5879 (2008).
- Heinen, A, et al. *Proc. Natl. Acad. Sci.* 105:8748 (2008).
- Bhattacharyya, BJ, et al. *J. Neurosci.* 28:6720 (2008).
- Samuels, IS, et al. *J. Neurosci.* 28:6983 (2008).
- Thonhoff, JR, et al., *Brain Res.* 1187:42 (2008).
- Watase, K, et al. *Proc. Natl. Acad. Sci.* 105:11987 (2008).
- Mitra, S, et al. *J. Biol. Chem.* 284:4398 (2009).
- Kohyama, J, et al. *Proc. Natl. Acad. Sci.* 105:18012 (2008).
- Schuetz, EG, et al. *J. Cell Physiol.* 134:309 (1988).
- DiPersio, CM, et al. *Mol. Cell Biol.* 11:4405 (1991).
- Mann, DJ, et al. *J. Mol. Endocrinol.* 8:235 (1992).
- Schuetz, JD and Scheutz EG, *Cell Growth and Diff.* 4:31 (1993).
- Rana, B, et al. *Mol. Cell Biol.* 14:5858 (1994).
- Lee, P, et al. *Proc. Natl. Acad. Sci.* 101:9263 (2004).
- Bi, Y, et al. *Drug Metab. Dispos.* 34:1658 (2006).
- Fahmi, OA, et al. *Drug Metab. Dispos.* 36:1971 (2008).
- Healan-Greenberg, C, et al. *Drug Metab. Dispos.* 36:500 (2008).

The above reference guide only represents a sample of citations for these products.

Product	Chemotaxis/Migration	Invasion	3D Culture	Neuronal Cells	Hepatocytes	Endothelial Cells	Tumor Cells	Muscle Cells	Epithelial Cells	Osteoclasts	hES and iPS cells	Non-specific Cell Attachment
	Fibronectin Is involved in cellular migration during wound healing and development as well as regulation of cell growth and differentiation.	3, 4, 7, 8, 10	11, 12	–	22	–	–	11, 65	59, 61, 62	–	–	76
Laminin Has been shown to stimulate neurite outgrowth, promote cell attachment, chemotaxis, cell differentiation, and neuronal survival.	9	–	–	22, 24, 26, 27, 29, 30, 31, 32, 34, 35, 37, 38, 39	–	–	64	–	–	–	–	–
Poly-D-Lysine Is a synthetic polymer used to promote attachment of a wide variety of cells, particularly neurons, glial cells, and transfected cells.	–	–	–	37, 38	–	–	–	–	–	–	–	–
Corning® PuraMatrix™ Peptide hydrogel Is a synthetic peptide that forms a clear 3D gel which mimics the fiber and pore size of the basement membrane. It can be used for in vitro and in vivo studies.	–	–	15, 20	21, 23, 36	15, 20	86	–	–	–	–	–	–
Vitronectin Mediates cell adhesion by integrin binding to the RGD (Arg-Gly-Asp) motif. Vitronectin participates in a variety of events including haemostasis, phagocytosis, tissue repair, and immune function.	6	–	–	–	–	–	–	–	–	71	–	–


Osteopontin is used to enhance attachment of tumor cell lines and osteoclasts. It is also a chemotactic factor for macrophages, smooth muscle cells, endothelial cells, and glial cells.

3D cell culture systems provide more in vivo-like culture conditions. Corning Matrigel® matrix, Corning Collagen I, Corning Laminin/Entactin high concentration, and Corning PuraMatrix Peptide hydrogel have been developed to meet the wide range of 3D culture needs, from reconstituted basement membrane to synthetic, defined surfaces.


49. Di Simone, N, et al. *J. Endocrinol.* 189:691. (2006).
50. Hawkins, BJ, et al. *Mol. Cell Biol.* 27:7582 (2007).
51. Ingram, DA, et al. *Stem Cells.* 25:297 (2007).
52. Kong, D, et al. *Cancer Res.* 67:3310 (2007).
53. Lee, S, et al. *Cell.* 130:691 (2007).
54. Michaud-Levesque, J, et al. *Carcinogenesis.* 28:280 (2007).
55. Nishiyama, K, et al. *J. Biol. Chem.* 282:17200 (2007).
56. Schmidt, M, et al. *Development.* 134:2913 (2007).
57. Takeda, Y, et al. *Blood.* 109:1524 (2007).
58. Dugas, JC, et al. *J. Neurosci.* 28:8294 (2008).
59. Hosseinkhani, M, et al. *Stem Cells.* 24:571 (2007).
60. Mendias CL, et al. *Proc. Natl. Acad. Sci.* 105:388 (2008).
61. Cohen-Karni, T, et al. *Proc. Natl. Acad. Sci.* 106:7309 (2009).
62. Mori, K, et al. *J. Biol. Chem.* 284:5067 (2009).
63. Park, CC, et al. *Cancer Res.* 66:1526 (2006).
64. Calabrese, C, et al. *Cancer Cell.* 11:69 (2007).
65. Clark, ES, et al. *Cancer Res.* 67:4227 (2007).
66. Itoh, M, et al. *Cancer Res.* 67:4759 (2007).
67. Wang, X, et al. *Cell.* 128:129 (2007).
68. Postovit, LM, et al. *Proc. Natl. Acad. Sci.* 105:4329 (2008).
69. Radjabi, AR, et al. *J. Biol. Chem.* 283:2822 (2008).
70. Senoo, M, et al. *Cell.* 129:523 (2007).
71. Zou, et al., *J. Biol. Chem.* 176:877 (2007).
72. Yang, D, et al. *Endocrinology.* 149:41728 (2008).
73. Xu, C, et al. *Nat. Biotechnol.* 19:971 (2001).
74. Amit, M, et al. *Biol. Reprod.* 70:837 (2004).
75. Ludwig, T, et al. *Nat. Methods.* 3:637 (2006).
76. Ludwig, T, et al. *Nat. Biotechnol.* 24:185 (2006).
77. Harb, N, et al. *PLoS ONE.* 3:e3001 (2008).
78. Takahashi, K, et al. *Cell.* 131:1 (2007).
79. Yu, J, et al. *Science.* 381:1917 (2007).
80. Duebel, J, et al. *Neuron.* 49:81 (2006).
81. Callen, E, et al. *Cell.* 130:63 (2007).
82. Calderon, B, et al. *Proc. Natl. Acad. Sci.* 105:6121 (2008).
83. Park, HS, et al. *J. Biol. Chem.* 283:26081 (2008).
84. Haggie, PM & Verkman, AS. *J. Biol. Chem.* 284:7681 (2009).
85. Skals, M, et al. *Proc. Natl. Acad. Sci.* 106:4030 (2009).
86. Sieminski, AL, et al. Primary sequence of ionic self-assembling peptide gels affects endothelial cell adhesion and capillary morphogenesis. *J. Biomed. Mater. Res. A* 87:494 (2008).

Corning® Extracellular Matrix Proteins

Get high quality data with Corning ECMs.



PC12 neurite outgrowth on Corning Collagen I.



Corning Gentest™ choly-lysyl-fluorescein sequestered in the bile canaliculi of Corning Gentest inducible-qualified human cryohepatocytes cultured on Corning BioCoat™ collagen I overlaid with Corning Matrigel matrix.

Description	Species	Qty.	Cat. No.
Corning Extracellular Matrix Proteins and Attachment Factors			
Corning Matrigel® matrix	mouse	5 mL	356234
	mouse	10 mL	354234
	mouse	50 mL (5 x 10 mL)	356235
Corning Matrigel matrix, high concentration	mouse	10 mL	354248
Corning Matrigel matrix, phenol red-free	mouse	10 mL	356237
Corning Matrigel matrix, growth factor reduced	mouse	5 mL	356230
	mouse	10 mL	354230
Corning Matrigel matrix, high concentration, growth factor reduced	mouse	10 mL	354263
Corning Matrigel matrix, phenol red-free, growth factor reduced	mouse	10 mL	356231
Corning Matrigel hESC-qualified matrix	mouse	5 mL	354277
Corning Cell-Tak™ cell and tissue adhesive	<i>Mytilus edulis</i>	1 mg	354240
	<i>Mytilus edulis</i>	5 mg	354241
	<i>Mytilus edulis</i>	10 mg (2 x 5 mg)	354242
Collagen I	bovine	30 mg	354231
	human	0.25 mg	354243
	human	10 mg	354265
	rat tail	100 mg	354236
	rat tail	1 g (10 x 100 mg)	356236
Collagen I, high concentration	rat tail	100 mg	354249
Collagen II	bovine	5 mg	354257
Collagen III	human	0.25 mg	354244
Collagen IV	human	0.25 mg	354245
	mouse	1 mg	354233
	mouse	10 mg (10 x 1 mg)	356233
Collagen V	human	0.25 mg	354246
Collagen VI	human	0.5 mg	354261
Fibronectin	human	1 mg	354008
	human	5 mg	356008
	human	25 mg (5 x 5 mg)	356009
Laminin	mouse	1 mg	354232
Ultra-pure laminin	mouse	1 mg	354239
Laminin/entactin complex, high concentration	mouse	10 mg	354259
Osteopontin	human	50 µg	354256
Poly-D-Lysine	–	20 mg	354210
Corning PuraMatrix™ peptide hydrogel	–	5 mL	354250
Vitronectin	human	0.25 mg	354238

Corning acquired the Discovery Labware Business including the BioCoat™, Gentest™, and Matrigel® brands. For information, visit www.corning.com/discoverylabware.

For Research Use Only. Not intended for use in diagnostic or therapeutic procedures.

For a listing of trademarks, visit us at www.corning.com/lifesciences/trademarks.

All other trademarks are property of their respective owners.

Corning Incorporated, One Riverfront Plaza, Corning, NY 14831-0001

CORNING

Corning Incorporated
Life Sciences

Corning BV
Fogstraat 12
1060 LJ Amsterdam
The Netherlands
Phone: +31 0 20 659 60 51
Fax: +31 20 659 76 73
cceurnl@corning.com

www.corning.com/lifesciences